


# RISK FACTORS FOR EQUINE LAMINITIS

Factors which make laminitis more likely to occur – CARE study findings<sup>1</sup>




High risk horses & ponies requiring extra vigilance




## Study Background



1,070 horses/ponies in Great Britain



Regular updates on health & management over 29 months



123 laminitis episodes reported

## References

<sup>1</sup> Pollard, D., Wylie, C.E., Verheyen, K.L.P. and Newton, J.R. (2019) Identification of modifiable factors associated with owner-reported equine laminitis in Britain using a web-based cohort study approach. BMC Veterinary Research 15:59 <https://doi.org/10.1186/s12917-019-1798-8>  
<sup>2</sup> Wylie, C.E., Collins, S.N., Verheyen, K.L., Newton, J.R., 2013b. Risk factors for equine laminitis: a case-control study conducted in veterinary-registered horses and ponies in Great Britain between 2009 and 2011. Vet. J. 198, 57-69 <https://doi.org/10.1016/j.tvjl.2013.08.028>

## Acknowledgements

Thanks go to all participants and contributors. Funding support: World Horse Welfare and the Margaret Giffen Charitable Trust. Collaborators: Animal Health Trust, Royal Veterinary College and Rossdale Equine Hospital.

## More information

✉ [danica.pollard@aht.org.uk](mailto:danica.pollard@aht.org.uk)

📘 [facebook.com/CAREstudy](https://www.facebook.com/CAREstudy)

[www.aht.org.uk/research/lameness/laminitis](http://www.aht.org.uk/research/lameness/laminitis)